

REVERSING SIGNS OF HAIR AGING:

Clinical Evidence of Gray Hair Pigmentation and Hair Quality Improvement

Layane Melody da Silva Reis Cavalcante¹, Lucia de Fatima Holanda¹, Giulia Okamoto Maciente¹, Mauricio Frazilho¹, Carla De Figueiredo Presti Favaro¹, Marisa Sacuragui¹, **Ana Lúcia Tabarini Alves Pinheiro**², Carla Monserrat Grecco Lopes², Mayara Miranda Ferreira², Viviane Cristina Albarici², Juliana Rodrigues Pinto²

¹Mantecorp Skin Care, Hynova, São Paulo/SP – Brazil
²Kosmoscience Group, Valinhos/SP – Brazil

P0463
EADV
2026



BACKGROUND

Hair graying is a prominent sign of aging, characterized by progressive loss of hair pigmentation, primarily associated with melanocyte dysfunction and influenced by multiple genetic, physiological, environmental, lifestyle, nutritional, and medical factors. Beyond its aesthetic impact, gray hair may affect self-image, confidence, and psychological well-being, driving demand for effective, non-invasive cosmetic strategies to restore a more youthful appearance while supporting scalp and hair health.

OBJECTIVE

To evaluate gray hair color recovery by image analysis and perceived improvements in hair health and quality following 90 days of use of **SHAMPOO DC-21051-33** and **LOTION DC-21052-06**.

STUDY DESIGN




Research Ethics Committee CAAE
68991223.1.0000.5514



17 participants
Mean age:
49 ± 7 years



90 days of home use
Evaluations:
Baseline
45 days
90 days




Shampoo DC-21051-33
and
Lotion DC-21052-06




Color Recovery by Image Analysis
ROI → Color histogram (C) →
Lower C = less gray hair
Perceived efficacy
5-point Likert scale
(1= strongly disagree; 5= strongly agree)


FORMULATION ACTIVES



BIOTIN




NANOENCAPSULATED TRICHOGENIC AUXIN



BLEND OF AMINO ACIDS

SELF-PERCEIVED IMPROVEMENTS (%)



All attributes achieved acceptance rates ≥ 70% after both 45 and 90 days of use.


OBJECTIVE EFFICACY – GRAY HAIR COLOR RECOVERY

AFTER 90 DAYS OF USE



13.7 – 20.4%

REDUCTION IN GRAY HAIR COLOR VARIATION ($p < 0.05$)



71% of participants showed recovery of gray hair coloration


REPRESENTATIVE CLINICAL IMAGES



Visual improvement in hair color uniformity and reduction of gray/white appearance.

INSTRUMENTAL IMAGE ANALYSIS

Gray Scale Histogram_Participant 07



The shift in gray-scale distribution after treatment indicates a measurable change in hair color toward darker tones.

CONCLUSION

After 45 and 90 days of combined use, SHAMPOO DC-21051-33 and LOTION DC-21052-06 showed high perceived efficacy across all evaluated hair health, growth, and pigmentation attributes.
After 90 days of combined use, objective image analysis demonstrated evidence of gray hair color recovery.
The combination represents a promising non-invasive cosmetic approach to support hair pigmentation and overall hair quality.

REFERENCES

- 1- Herdiana Y. Gray hair: from preventive to treatment. *Clin Cosmet Investig Dermatol*. 2025;18:1475-1494. doi:10.2147/CCID.S526263
- 2 - Saad M, Babar NF, Majeed R, et al. Impact of premature greying of hair on socio-cultural adjustment and self-esteem among medical undergraduates in Foundation University, Islamabad. *Cureus*.2019;11(7):e5083.doi:10.7759/cureus.5083.
- 3 - Santos AC, Panchal A, Rahman N, Pereira-Silva M, Pereira I, Veiga F, Lvov Y. Evolution of hair treatment and care: prospects of nanotube-based formulations. *Nanomaterials (Basel)*. 2019;9(6):903. doi:10.3390/nano9060903.
- 4- Mirmirani P. Age-related hair changes in men: mechanisms and management of alopecia and graying. *Maturitas*. 2015;80(1):58-62. doi:10.1016/j.maturitas.2014.10.008.